

# Covid-19 Vaccine

## Who should get a booster?

The CDC recommends people age 5 and older who received Pfizer and 18 and older who received Moderna or Johnson & Johnson get a booster dose when they are eligible: five months after their second dose of a Moderna or Pfizer vaccine and two months after a Johnson & Johnson (Janssen) vaccine dose. People 5 and older who received Pfizer or 18 and older who received Moderna who are immunocompromised and received a third dose are recommended to receive a booster dose three months after their third dose. People who are 50 years of age and older, immunocompromised persons at least 12 years old who received Pfizer or at least 18 years old and received Moderna or Johnson & Johnson, and persons who received both a primary and booster dose of Johnson & Johnson should receive a second booster dose of an mRNA vaccine four months after their first booster dose.

### Moderna:

- Third dose of Moderna is recommended for immunocompromised people age 6 months and older at least 28 days after the second dose.
- This additional dose is a part of the primary vaccine series. Immunocompromised people 18 and older should receive a booster dose three months after their additional dose and a second booster dose four months after their first booster dose.

### Pfizer:

- Third dose of Pfizer is recommended for immunocompromised people age 5 and older at least 28 days after the second dose.
- This additional dose is a part of the primary vaccine series. A booster dose is recommended for immunocompromised people 5 and older three months or more after their third dose.

### Johnson & Johnson

- For people ages 18 and older who are immunocompromised and who received Johnson & Johnson (Janssen) vaccine, a second dose of an mRNA COVID-19 vaccine (Moderna or Pfizer) is recommended at least 28 days after the first dose.
- Everyone who received a Johnson & Johnson (Janssen) vaccine is advised to get a booster dose. An mRNA vaccine is recommended.

People who are immunocompromised may receive a second booster dose of an mRNA vaccine four months after their first booster dose or four months after their additional dose (if the additional dose was received after the booster dose.)

